



A very large (effectively infinite) capacitor has charge Q . A neutral (*homogeneous*) dielectric is inserted into the gap (and of course, it will polarize). We want to find \mathbf{E} everywhere.



Which equation would you head to first?

- A. $\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$
- B. $\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$
- C. $\oint \mathbf{E} \cdot d\mathbf{A} = \frac{Q}{\epsilon_0}$
- D. More than one of these would work
- E. Can't solve unless we know the dielectric is linear.


A very large (effectively infinite) capacitor has charge Q . A neutral (*homogeneous*) dielectric is inserted into the gap (and of course, it will polarize). We want to find \mathbf{D} everywhere.



Which equation would you head to first?

- A. $\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$
- B. $\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$
- C. $\oint \mathbf{E} \cdot d\mathbf{A} = \frac{Q}{\epsilon_0}$
- D. More than one of these would work

An ideal (large) capacitor has charge Q . A neutral linear dielectric is inserted into the gap. We want to find \mathbf{D} in the dielectric.




$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

For the Gaussian pillbox shown, what is $Q_{free,enclosed}$?

- A. σA
- B. $-\sigma_B A$
- C. $(\sigma - \sigma_B)A$
- D. $(\sigma + \sigma_B)A$
- E. Something else

An ideal (large) capacitor has charge Q . A neutral linear dielectric is inserted into the gap. We want to find \mathbf{D} in the dielectric.




$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

Is \mathbf{D} zero INSIDE the metal? (i.e., on the top face of our cubical Gaussian surface)

- A. It must be zero in there.
- B. It depends.
- C. It is definitely not zero in there.

An ideal (large) capacitor has charge Q . A neutral linear dielectric is inserted into the gap. We want to find \mathbf{D} in the dielectric.




$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

What is $|\mathbf{D}|$ in the dielectric?


- A. σ
- B. 2σ
- C. $\sigma/2$
- D. $\sigma + \sigma_b$
- E. Something else

An ideal (large) capacitor has charge Q . A neutral linear dielectric is inserted into the gap (with given dielectric constant). Now that we have \mathbf{D} in the dielectric, what is \mathbf{E} in that **small gap** above the dielectric?




- A. $\mathbf{E} = \mathbf{D}\epsilon_0\epsilon_r$
- B. $\mathbf{E} = \mathbf{D}/\epsilon_0\epsilon_r$
- C. $\mathbf{E} = \mathbf{D}\epsilon_0$
- D. $\mathbf{E} = \mathbf{D}/\epsilon_0$
- E. Not so simple! Need another method

An ideal (large) capacitor has charge Q . A neutral linear dielectric is inserted into the gap. Now that we have \mathbf{D} in the dielectric, what is \mathbf{E} inside the dielectric?



- A. $\mathbf{E} = \mathbf{D}\epsilon_0\epsilon_r$
- B. $\mathbf{E} = \mathbf{D}/\epsilon_0\epsilon_r$
- C. $\mathbf{E} = \mathbf{D}\epsilon_0$
- D. $\mathbf{E} = \mathbf{D}/\epsilon_0$
- E. Not so simple! Need another method


An ideal (large) capacitor has charge Q . A neutral linear dielectric is inserted into the gap (with given dielectric constant). Where is \mathbf{E} discontinuous?



- i) near the free charges on the plates
- ii) near the bound charges on the dielectric surface

- A. i only
- B. ii only
- C. both i and ii (but nowhere else)
- D. both i and ii but also other places
- E. none of these/something else

An ideal (large) capacitor has charge Q . A neutral linear dielectric is inserted into the gap (with given dielectric constant).




Where is \mathbf{E} discontinuous?

- i) near the free charges on the plates
- ii) near the bound charges on the dielectric surface

- A. i only
- B. ii only
- C. both i and ii (but nowhere else)
- D. both i and ii but also other places
- E. none of these/something else

An ideal (large) capacitor has charge Q . A neutral linear dielectric is inserted into the gap (with given dielectric constant).




Where is \mathbf{D} discontinuous?

- i) near the free charges on the plates
- ii) near the bound charges on the dielectric surface

- A. i only
- B. ii only
- C. both i and ii (but nowhere else)
- D. both i and ii but also other places
- E. none of these/something else


A point charge $+q$ is placed at the center of a neutral, linear, homogeneous, dielectric teflon shell. Can \mathbf{D} be computed from its divergence?



$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

- A. Yes
- B. No
- C. Depends on other things not given


A point charge $+q$ is placed at the center of a neutral, linear, homogeneous dielectric teflon shell. The shell polarizes due to the point charge. Is the curl of the polarization \mathbf{P} zero everywhere?



$$\oint \mathbf{P} \cdot d\mathbf{l} = 0 \text{ for every loop?}$$

- A. Yes
- B. No
- C. Depends on other things not given


A point charge $+q$ is placed at the center of a neutral, linear, dielectric **hemispherical** shell. Can \mathbf{D} be computed from its divergence?



- A. Yes
- B. No
- C. Depends on the inner radius of the dielectric

$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

A point charge $+q$ is placed at the center of a neutral, linear, dielectric shell. The shell polarizes due to the point charge. Is the curl of the polarization \mathbf{P} zero everywhere?



- A. Yes
- B. No
- C. Depends on the inner radius of the dielectric.

$$\oint \mathbf{P} \cdot d\mathbf{l} = 0 \text{ for every loop?}$$