

What is the physical interpretation of $\oint \mathbf{A} \cdot d\mathbf{l}$?

- A. The current density \mathbf{J}
- B. The magnetic field \mathbf{B}
- C. The magnetic flux Φ_B
- D. It's none of the above, but is something simple and concrete
- E. It has no particular physical interpretation at all

Consider a square loop enclosing some amount of magnetic field lines with height H and length L . We intend to compute $\Phi_B = \oint \mathbf{A} \cdot d\mathbf{l}$? What happens to Φ_B as H becomes vanishingly small?

- A. Φ_B stays constant
- B. Φ_B gets smaller but doesn't vanish
- C. $\Phi_B \rightarrow 0$

Consider a square loop enclosing some amount of magnetic field lines with height H and length L . If $\Phi_B \rightarrow 0$ as $H \rightarrow 0$ (or $L \rightarrow 0$), what does that say about the continuity of \mathbf{A} ?

$$\Phi_B = \oint \mathbf{A} \cdot d\mathbf{l}$$




- A. \mathbf{A} is continuous at boundaries
- B. \mathbf{A} is discontinuous at boundaries
- C. ???

The leading term in the vector potential multipole expansion involves:

$$\oint d\mathbf{l}'$$

What is the magnitude of this integral?

- A. R
- B. $2\pi R$
- C. 0
- D. Something entirely different/it depends!



Two magnetic dipoles m_1 and m_2 (equal in magnitude) are oriented in three different ways.

Which ways produce a dipole field at large distances?

- A. None of these
- B. All three
- C. 1 only
- D. 1 and 2 only
- E. 1 and 3 only