



5 charges,  $q$ , are arranged in a regular pentagon, as shown.  
What is the E field at the center?



- A. Zero
- B. Non-zero
- C. Really need trig and a calculator to decide

1 of the 5 charges has been removed, as shown.  
What's the E field at the center?



- A.  $+(kq/a^2)\hat{y}$
- B.  $-(kq/a^2)\hat{y}$
- C. 0
- D. Something entirely different!
- E. This is a nasty problem which I need more time to solve

## ANNOUNCEMENTS


- Help Session 1420 BPS (4-5pm)
  - Starts this week!
- We will use GitHub Classroom for [digital submissions of homework](#)
  - Create a [GitHub account](#)
  - Download [GitHub Desktop](#)
  - Review [Piazza post on usage](#)
  - Come to help session (or my office) if you need/want help

To find the E-field at P from a thin line (uniform charge density  $\lambda$ ):

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\lambda d\mathbf{l}'}{\mathcal{R}^2} \hat{\mathcal{R}}$$

What is  $\mathcal{R}$ ?

- A.  $x$
- B.  $y'$
- C.  $\sqrt{dl'^2 + x^2}$
- D.  $\sqrt{x^2 + y'^2}$
- E. Something else



What do you expect to happen to the field as you get really far from the rod?

$$E_x = \frac{\lambda}{4\pi\epsilon_0} \frac{L}{x\sqrt{x^2 + L^2}}$$

- A.  $E_x$  goes to 0.
- B.  $E_x$  begins to look like a point charge.
- C.  $E_x$  goes to  $\infty$ .
- D. More than one of these is true.
- E. I can't tell what should happen to  $E_x$ .




Which of the following are vectors?

(I) Electric field, (II) Electric flux, and/or (III) Electric charge



- A. I only
- B. I and II only
- C. I and III only
- D. II and III only
- E. I, II, and III

Given the location of the little bit of charge ( $dq$ ), what is  $|\vec{R}|$ ?



- A.  $\sqrt{z^2 + r'^2}$
- B.  $\sqrt{z^2 + r'^2 - 2zr' \cos \theta}$
- C.  $\sqrt{z^2 + r'^2 + 2zr' \cos \theta}$
- D. Something else


A positive point charge  $+q$  is placed outside a closed cylindrical surface as shown. The closed surface consists of the flat end caps (labeled A and B) and the curved side surface (C). What is the sign of the electric flux through surface C?




- A. positive
- B. negative
- C. zero

D. not enough information given to decide

Let's get a better look at the side view.



Which of the following two fields has zero divergence?



- A. Both do.
- B. Only I is zero
- C. Only II is zero
- D. Neither is zero
- E. ???


What is the value of:

$$\int_{-\infty}^{\infty} x^2 \delta(x - 2) dx$$


- A. 0
- B. 2
- C. 4
- D.  $\infty$
- E. Something else

A point charge ( $q$ ) is located at position  $\mathbf{R}$ , as shown. What is  $\rho(\mathbf{r})$ , the charge density in all space?

- A.  $\rho(\mathbf{r}) = q\delta^3(\mathbf{R})$
- B.  $\rho(\mathbf{r}) = q\delta^3(\mathbf{r})$
- C.  $\rho(\mathbf{r}) = q\delta^3(\mathbf{R} - \mathbf{r})$
- D.  $\rho(\mathbf{r}) = q\delta^3(\mathbf{r} - \mathbf{R})$
- E. Something else??




An electric dipole (+ $q$  and  $-q$ , small distance  $d$  apart) sits centered in a Gaussian sphere.



What can you say about the flux of  $\mathbf{E}$  through the sphere, and  $|\mathbf{E}|$  on the sphere?

- A. Flux = 0,  $E = 0$  everywhere on sphere surface
- B. Flux = 0,  $E$  need not be zero *everywhere* on sphere
- C. Flux is not zero,  $E = 0$  everywhere on sphere
- D. Flux is not zero,  $E$  need not be zero...

Which of the following two fields has zero curl?




- A. Both do.
- B. Only I is zero
- C. Only II is zero
- D. Neither is zero
- E. ???

Can superposition be applied to electric potential,  $V$ ?

$$V_{tot} \stackrel{?}{=} \sum_i V_i = V_1 + V_2 + V_3 + \dots$$




- A. Yes
- B. No
- C. Sometimes



Could this be a plot of  $|\mathbf{E}(r)|$ ? Or  $V(r)$ ? (for SOME physical situation?)


- A. Could be  $E(r)$ , or  $V(r)$
- B. Could be  $E(r)$ , but can't be  $V(r)$
- C. Can't be  $E(r)$ , could be  $V(r)$
- D. Can't be either
- E. ???

A point charge  $+q$  sits outside a **solid neutral conducting copper sphere** of radius  $A$ . The charge  $q$  is a distance  $r > A$  from the center, on the right side. What is the E-field at the center of the sphere? (Assume equilibrium situation).



- A.  $|E| = kq/r^2$ , to left
- B.  $kq/r^2 > |E| > 0$ , to left
- C.  $|E| > 0$ , to right
- D.  $E = 0$
- E. None of these

A neutral copper sphere has a spherical hollow in the center. A charge  $+q$  is placed in the center of the hollow. What is the total charge on the outside surface of the copper sphere? (Assume Electrostatic equilibrium.)



- A. Zero
- B.  $-q$
- C.  $+q$
- D.  $0 < q_{outer} < +q$
- E.  $-q < q_{outer} < 0$