

In our basic model of a battery, we said that the force per unit charge the battery provides through chemical reactions (\mathbf{f}) had a magnitude equal the electric field produced by the separated charge (\mathbf{E}). This maintains a constant drift speed across the battery for the electrons.


If the battery has some internal resistance (r), which is larger?

- A. $|\mathbf{f}|$
- B. $|\mathbf{E}|$
- C. Both still the same

A metal bar moves with constant speed **to the right**. A constant magnetic field points **out of the page**. What happens to the electrons in the bar (in the frame of the moving bar)?


- A. Nothing
- B. They move upward
- C. They move downward
- D. They move left
- E. They move right

One end of rectangular metal loop enters a region of constant uniform magnetic field \mathbf{B} , with initial constant speed v , as shown. What direction is the magnetic force on the loop?




- A. Up the "screen" \uparrow
- B. Down the "screen" \downarrow
- C. To the right \rightarrow
- D. To the left \leftarrow
- E. The net force is zero

One end of rectangular metal loop enters a region of constant uniform magnetic field \mathbf{B} , out of page, with constant speed v , as shown. As the loop enters the field is there a non-zero emf around the loop?



- A. Yes, current will flow CW
- B. Yes, current will flow CCW
- C. No

A rectangular metal loop moves through a region of constant uniform magnetic field \mathbf{B} , with speed v at $t = 0$, as shown. What is the magnetic force on the loop at the instant shown? Assume the loop has resistance R .



A. $2L^2 v B^2 / R$ (right)

B. $2L^2 v B^2 / R$ (left)

C. 0

D. Something else/not sure...